



## **STANDARDS FOR INTERVENTIONAL ELECTROPHYSIOLOGY STUDY AND CATHETER ABLATION IN ADULTS – February 2016**

### **1. INTRODUCTION**

This document replaces the previous British Heart Rhythm Society (BHRS) document “Standards for interventional electrophysiology study and catheter ablation in adults” issued in January 2014. It has been compiled by a group of interventional cardiac electrophysiologists, cardiac physiologists and specialist arrhythmia nurses drawn from both secondary and tertiary centres and approved by the Council of the BHRS in February 2016. This document will be reviewed by BHRS Council no later than February 2018.

The purpose of the document is to facilitate the provision of high quality, evidence-based electrophysiology (EP) assessment and intervention to all patients who may benefit. This includes the identification of patients with indications for intervention, a broad description of the range of procedures available and the means by which to provide them, patient and procedure follow up, data collection, storage and submission.

It includes the best available evidence and expert opinion on current practice. The source material for this evidence is listed in the reference section.

It is recognised that competency can only be defined effectively in terms of patient outcome. Numbers given in this document are indicative and should not be taken in isolation as evidence of competency or the ability to provide a safe, high quality service. This document is not intended to disrupt or disenfranchise existing, successful electrophysiology services. It should be regarded as a template for developing best practice when starting *de novo* and a recommendation to enable successful but inadequately resourced services to develop.

This document is not intended to replace Trust policies and other legislation e.g. data protection and codes of conduct that should be adhered to in addition to the recommendations of this document.

In summary, this document aims to:

- Deliver the highest possible standard of care for patients with arrhythmias
- Encourage strong links and active collaboration between primary care, secondary care EP units and tertiary centre EP units
- Ensure that procedures are performed by appropriately skilled operators in a suitably staffed and equipped environment achieving nationally acceptable success rates with minimised complication rates
- Ensure that patients have 24/7 access to specialist arrhythmia care and advice at regional level

## 2. DEFINITIONS

The following definitions are used within this document. For the purposes of this document, as some Trusts cover multiple sites, an electrophysiology centre is taken to mean a single hospital site where invasive electrophysiology procedures are performed rather than the Trust as a whole. It is accepted that operators may work at more than one centre but each centre should conform to the standards within this document. Furthermore, as more non-cardiac surgical centres develop in-house interventional electrophysiology capabilities, it is appropriate to describe the scope of practice as *standard* catheter ablation and *complex* catheter ablation. BHRS recognises the difficulty in defining absolute requirements for this artificial distinction between intervention types and acknowledges that locally available systems of care and expertise will contribute critically to the scope of locally available electrophysiology practice.

### ***Standard EP study / catheter ablation:***

Invasive electrophysiology study

Ablation of the atrioventricular node

Ablation of atrioventricular nodal reentrant tachycardia

Ablation of accessory pathways

Ablation of cavo-tricuspid isthmus dependent atrial flutter

### ***Complex catheter ablation:***

Ablation of atrial fibrillation (AF)

Ablation of non-cavotricuspid isthmus-dependent atrial tachycardia

Ablation of ventricular tachycardia (VT) including idiopathic focal ventricular tachycardia

Ablation of atrial or ventricular arrhythmias via the pericardial route

Modification of the sinus node

Ablation of in patients with complex congenital heart disease (see BHRS document "Cardiac Rhythm management in patients with congenital heart disease").

### ***First line treatment:***

The therapy is offered as the treatment of first choice

### ***Second line treatment:***

The therapy is offered after the patient has found other treatments unsuccessful or intolerable (antiarrhythmic medication, cardiac implantable devices)

### ***Operator:***

The first operator is the clinician who is the main individual performing the case, especially at the time of the ablation. The first operator will normally be scrubbed, with the exception of some cases where the majority of the procedure is conducted robotically. A second operator is another clinician present throughout and normally also scrubbed. Either first or second operator can be an accredited electrophysiologist or a trainee. The responsible consultant should be an accredited electrophysiologist supervising the case, who can also be one of the actual operators. For the purpose of recording procedure numbers, a clinician can "count" procedures conducted in any of these three roles, though those where he/she was not physically operating should be enumerated separately.

### **Training Centres:**

At present, there is no mechanism for designation of Training Centres in the UK although it is likely that a process of departmental accreditation in Training and Practice will be developed in the future. However, the Joint Royal College of Physicians Training Board (JRCPTB) Speciality Training Curriculum for Cardiologists (August 2010)<sup>1</sup> specifies that centres offering advanced rhythm training should have at least 2 electrophysiologists who spend the majority of their time practising electrophysiology. Centres should perform  $\geq 100$  EP studies and catheter ablations per year and have availability of complex mapping and ablation technologies.

## **3. TREATMENT INDICATIONS**

Catheter ablation is primarily a therapy that improves quality of life although in certain situations may improve prognosis. A total of 17,578 ablation procedures were performed in the United Kingdom in 2014 across 61 centres with a mean number of 288 ablations per centre. Thirty-six centres performed  $>50$  AF ablations and 23 centres performed  $>10$  structural heart VT ablations.<sup>4</sup> The various indications for catheter ablation are clearly described in numerous international guidelines and consensus statements<sup>5-7</sup> and the National Service Framework for Coronary Heart Disease chapter 8 with some aspects of catheter ablation already examined and approved by NICE.<sup>8-10</sup> It is not the purpose of this document to reinterpret peer reviewed, published guidance. It is recognised, however, that such guidance may not cover all patient groups and may not be appropriate in certain situations. Furthermore, clinical judgement based on published evidence should be used for indications not yet considered by NICE. It is important to demonstrate compliance with best practice and regular audit of procedure indications and outcomes is strongly recommended. It is essential that all interventional EP centres ensure accurate and timely procedure data submission to the national CRM database.

In summary, BHRS recommends that catheter ablation is offered as:

### **First line treatment for:**

All regular supraventricular tachycardias (SVTs) including cavo-tricuspid isthmus dependent atrial flutter that are either causing symptoms or have potential to produce tachycardia-related cardiomyopathy

### **Second line treatment for:**

Symptomatic AF

AF (symptomatic or asymptomatic) that has potential to produce or be associated with tachycardia cardiomyopathy (rate control with AV node ablation and pacing or rhythm control with AF ablation)

Symptomatic VT, VT storm or VT leading to intolerable or very frequent ICD therapy

VT with the potential to produce tachycardia-related cardiomyopathy

For complex ablation, the development of a multidisciplinary approach to patient selection, management and follow up is recommended. Involvement of physicians (interventional electrophysiologists, imaging and heart failure specialists),

physiologists, and specialist arrhythmia nurses to aid in assessment of suitability for ablation and follow up may lead to improved patient outcomes. BHRS recommends that the appropriate international guidelines be used to inform practice and that the reasons for practice outside well established and accepted guidelines and consensus statements be clearly justified.

#### **4. REQUIREMENTS FOR PERFORMING CATHETER ABLATION**

Safe invasive electrophysiology assessment and treatment requires the appropriate environment, equipment, trained personnel and culture.<sup>11</sup> BHRS recommends that all centres and cardiologists performing EP studies should also have competency to proceed safely to catheter ablation if necessary including access to the left atrium and great vessels if required. All electrophysiology studies and ablations should be performed in units that meet or exceed the quality standards, with a minimum of two active consultant cardiac electrophysiologists per centre, one of whom should ideally hold BHRS, EHRA or IBHRE certification. Formal arrangements for handover, in-hours and out-of-hours expert cover and on-going care should be in place.

Cardiac arrhythmias are a major source of morbidity and mortality for patients with Adult Congenital Heart Disease (ACHD). BHRS has produced a separate standards document for cardiac rhythm management in these patients. BHRS recognises that advanced interventional EP training in ACHD is unlikely to be achieved within the 2 year subspecialty training programme and should be the focus of further training for appropriate individuals. These patients are often complex and their care should be coordinated through a multidisciplinary meeting including specialist electrophysiology input. Electrophysiology interventional procedures in ACHD patients should include an adult cardiac electrophysiologist trained in complex adult EP intervention. More in depth cross-training of electrophysiologists from both paediatrics and adult cardiology is needed to meet the special needs of these patients. It is reasonable to recommend that interventional arrhythmia procedures be performed at centres where the staff are experienced with the complex anatomy and distinctive arrhythmia substrates encountered in congenital heart defects.<sup>12</sup> These procedures should not be performed in non-specialist centres or by individuals without the appropriate expertise in cardiac electrophysiology.

##### **4.1 CARDIOLOGISTS – training requirements**

This section refers to cardiologists performing standard and complex catheter ablation procedures.

- a. Trainees should have achieved a CCT in cardiology and completed a recognised EP training program in a Deanery-designated training centre or equivalent. The programme should include at least 2 years of subspecialty training in heart rhythm management with dedicated time within that programme specialising in electrophysiology and ablation
- b. Trainees should have completed a log book of procedures
- c. Trainees should have been actively involved with at least 200 standard ablation procedures and performed 50 as first operator
- d. Trainees should have completed training in transseptal puncture and performed >20 transseptal punctures as first operator

- e. Trainees may also elect to train in more complex ablation techniques, the requirements for which are more exacting than for standard ablation.<sup>13</sup> To achieve the minimum standard for training, trainees should have performed:
  - For complex atrial arrhythmias, to have been actively involved with at least 100 complex atrial ablation (primarily AF and left atrial tachycardia) with 50 as first operator
  - For ventricular arrhythmias in structural heart disease or involving 3D electroanatomic mapping – been actively involved with 20 ablations with 10 as first operator
  - For complex congenital heart disease – been actively involved with 20 ablations with 10 as first operator
  - For epicardial ablation, been actively involved with 10 “dry” pericardial punctures and to have performed 5 “dry” punctures as the first operator
- f. Trainees should not be permitted to perform procedures unsupervised until they have received a certificate of completion of specialist training. BHRS recognises that the level of supervision will appropriately vary according to the expertise of the individual trainee
- g. Trainers should be in whole time employment and spend at least 70% of their working time involved in heart rhythm management. Cardiac electrophysiologists who are working part time can contribute to training but should not be the primary clinical or education supervisor.
- h. Trainers should be actively involved in heart rhythm audit and research
- i. Trainers should achieve the standards required for maintenance of competence
- j. Trainers should be aware of the recommendations by the professional bodies pertaining to the specialty, have annual appraisal and maintain a CPD diary

#### 4.2 CARDIOLOGISTS – maintenance of competence

- a. There should be a minimum of 2 active interventional cardiac electrophysiologists per centre<sup>13</sup> (one of whom should ideally be BHRS, EHRA or IBHRE certified), with formal support from larger centres encouraged for centres establishing de novo invasive EP services. Although it is possible to perform catheter ablation in a centre with one competent specialist, this is not a sustainable model of care for the development of a high quality, comprehensive heart rhythm service
- b. Complex ablation should not be performed in single operator centres
- c. All interventional electrophysiologists should perform at least 50 catheter ablation procedures per year as an “operator” (as defined above). For those undertaking complex ablation, a minimum of 25 procedures should be complex, however, 50 is desirable. Those undertaking less than 25 AF ablations per year have a demonstrably higher complication rate.<sup>14</sup> An operator performing more than 50 complex ablations per year will be deemed to have met the minimum requirements for simple and complex ablations. The association between complications and case volume is acknowledged by BHRS and is the justification behind these minimum basic requirements.<sup>14</sup> BHRS recognises the changing case mix encountered during both training and consultant practice and encourages

all electrophysiologists to maintain their skills and knowledge in both simple and complex cases.<sup>15</sup>

- d. If a cardiologist does not perform this number of procedures per year then competence should be independently assessed in accordance with current specialty training guidelines\* (a structured report should be obtained from the trainer/assessor on competency for "sign off." This report should include the number of cases performed and the level of competency as assessed by the trainer/assessor. Assessment should consist of successful completion of 6 DOPS in catheter ablation at level 3 by at least 2 assessors)
- e. All cardiologists performing catheter ablation must undertake appropriate CPD in electrophysiology including implications for driving
- f. All electrophysiologists must audit their personal complications and share these within their centre at least quarterly and through the national CRM database for clinical governance purposes. If an operator's complications were to exceed accepted limits<sup>15</sup>, practice should be reviewed locally by the appropriate clinical governance structures already in place and if necessary, advice sought from elsewhere within the UK. Operators performing fewer than 50 procedures per year may need to average their figures over 2 or more years to account for random variation. For consultants engaged in a supervising role, complications occurring during a procedure will be ascribed to the consultant responsible for the patient for the purposes of data returns.

#### 4.3 CARDIAC PHYSIOLOGISTS / HEALTH CARE SCIENTISTS (HCSs)

- a. There should be at least 2 cardiac physiologists / HCSs actively involved in interventional electrophysiology where these procedures are performed
- b. Each physiologist / HCS must have had appropriate training in the execution of electrophysiology studies and catheter ablation
- c. At least 1 physiologist / HCS should have current accreditation in catheter ablation and electrophysiology (BHRS, EHRA or IBHRE)
- d. All physiologists / HCSs must undertake appropriate continuing professional development in catheter ablation and resources made available for them to do so
- e. Each physiologist / HCS should be actively involved in at least 30 electrophysiology procedures per year. If a physiologist / HCS does not perform this number of procedures in a 12 month period then competence should be independently reassessed in line with local governance procedures.

#### 4.4 SPECIALIST ARRHYTHMIA NURSES

- a. Hospitals delivering specialist cardiac arrhythmia care are expected to develop the role of Cardiac Arrhythmia Nurse Specialist as an appropriate (and sustainable) level as recommended in the NSF Chapter 8<sup>8</sup>. Ideally, at least 2 nurses per centre should be denoted as specialist arrhythmia nurses to allow continuity of care during periods of absence and can be achieved if necessary by nurses taking up dual or part time roles

---

\* Operators performing fewer than 50 procedures per year may need to average their figures over 2 years to account for random variation, however, the necessity to do this should be seen as an exception

- b. Cardiac arrhythmia nurses should receive training appropriate to their involvement in the CRM team and should work according to protocols developed within their centre and nationally
- c. Nurses should undertake appropriate CPD in heart rhythm management and resources made available for them to do so. Nurses should be encouraged and supported to undertake BHRS certification
- d. Where nurses are running outpatient clinics independently, they should have same day direct telephone access to a consultant and the opportunity to meet with the supervising consultant at least fortnightly to discuss clinical cases.

#### 4.5 CENTRES

In order to comply with the requirements for individual operators, centres performing catheter ablation procedures should undertake a minimum of 100 ablations per year. Centres undertaking catheter ablation for AF should also perform a minimum of 50 AF ablations per year. This should be regarded as a minimum and for centres offering advanced training in cardiac rhythm management, it would be expected that they would undertake significantly more procedures.

##### *Basic requirements – standard ablation:*

Centres performing catheter ablation procedures must have a minimum set of equipment to safely carry out these procedures. These should include:

- a. Wards with bedside cardiac monitoring equipment including effective rhythm alarm systems and capable of storing 24 hours of continuous ECG data
- b. Nursing staff experienced in care and management of patients with cardiac arrhythmias
- c. Facilities for cardiopulmonary resuscitation, temporary and permanent pacing
- d. A catheter laboratory equipped with modern X-ray equipment and the appropriate patient and operator radiation monitoring systems in place
- e. Facilities for anaesthesia and ventilation within the laboratory in the event that they are needed
- f. A modern digital electrophysiology recording and storage system and haemodynamic monitoring; a programmable stimulator; a radiofrequency generator specifically for EP procedures
- g. Equipment for pericardiocentesis
- h. Echocardiography

##### *Additional requirements – complex ablation:*

For centres undertaking complex ablation procedures, BHRS advises that there should be on site access to emergency cardiothoracic surgery or arrangements in place for immediate transfer. While acknowledging that this will be rarely employed, as the scope increases of complex left atrial, ventricular and epicardial procedures, it is crucial that optimum patient safety measures are in place to manage rare but severe complications. Where not available on-site, an agreed written protocol must be in place with the local cardiac surgical centre and local ambulance service to provide emergency transfer and surgical cover. In addition, the time taken for a patient to thoracotomy should be of a similar order to that possible with on-

site surgical facilities where a surgical team is not on stand-by. Centres undertaking complex ablation should have ready access to the following:

- a. Electroanatomic mapping hardware
- b. Haemodynamic assist devices (intra-aortic balloon pump, mechanical assist devices) for VT ablation in structural heart disease
- c. Intensive Therapy Unit availability for planned high-risk cases

## **5. POST PROCEDURE FOLLOW UP STANDARDS**

### *Standard ablation*

All patients who undergo catheter ablation should be followed up at least once in a clinic supervised by the operator to allow audit of the outcomes of the ablation performed. Some centres may choose to do this using remote follow up but time must be set aside to talk to patients (even by phone), examine their follow up data (ECG/Holter arranged locally) and log the outcome in a database. Follow up must also be consistent with national database requirements.

### *Complex ablation*

BHRS endorses the 2012 HRS/EHRA/ECAS expert consensus statement on catheter and surgical ablation of atrial fibrillation.<sup>7</sup> Some points deserve special mention:

- a. ECG confirmation of the diagnosis of AF is mandatory prior to recommending catheter ablation as a treatment
- b. Cardiac imaging to assess cardiac structure and function is recommended prior to catheter ablation
- c. Anticoagulation is recommended following catheter ablation, tailored to the stroke risk of the patient rather than to the anticipated effect on frequency of atrial fibrillation. The CHA<sub>2</sub>S<sub>2</sub>VASc scoring system should be used in preference to the CHADS<sub>2</sub> score
- d. Following successful catheter ablation, patients should be followed up for at least 1 year in the outpatient setting
- e. Patient related outcome measures (PROMS) should be collected following catheter ablation of AF
- f. Patients who have undergone ablation should be provided with a contact number at the ablating centre and advised to report symptoms occurring within the first six weeks following ablation. Atrial oesophageal fistula is likely under-reported and fatal if not recognised and treated early.<sup>16</sup> Diagnostic delay following post-procedure presentation to a non-ablation centre can be reduced by early communication with the ablation centre. Protocols should be in place for the diagnosis and management of this rare but catastrophic complication. It is recommended that specific information relating to atrio-oesophageal fistula (such as that offered by Arrhythmia Alliance) is provided to patients following catheter ablation of AF.

## **6. AUDIT**

Complications associated with catheter ablation typically occur at the time of the procedure or shortly thereafter i.e. within six weeks. There are also occasional



advice and safety notices from manufacturers and the MHRA which necessitate timely action.

- a. All ablation centres must collect data on their patients, procedures and follow-up which is immediately available and facilitates audit. It is mandatory that all hospitals performing catheter ablation submit accurate and timely procedural data to the national CRM database. This is a national quality requirement and is audited by the Care Quality Commission
- b. In addition, the audit data should be presented and discussed in a multidisciplinary team meeting at least quarterly. Attendance should be mandatory for trainers and trainees alike.

## REFERENCES

1. Speciality Training Curriculum for Cardiology. Joint Royal College of Physicians Training Board August 2010
2. Merino JL, Arribas F, Botto GL, et al. Core curriculum for the heart rhythm specialist. *Europace*. 2009;11:iii1-iii26
3. Merino JL, Arribas F, Botto GL, et al. Core curriculum for the heart rhythm specialist: Executive summary. *Europace*. 2009;11:1381-1386
4. Raatikainen MJ, Arnar DO, Zeppenfeld K, et al. Statistics on the use of cardiac implantable electronic devices and electrophysiological procedures in the European Society of Cardiology countries: 2014 report from the European Heart Rhythm Association. *Europace* 2015;17 suppl 1:i1-75.
5. Aliot EM, Stevenson WG, Almendral-Garrote JM, et al. EHRA/HRS expert consensus on catheter ablation of ventricular arrhythmias: Developed in a partnership with the european heart rhythm association (EHRA), a registered branch of the european society of cardiology (ESC), and the heart rhythm society (HRS); in collaboration with the american college of cardiology (ACC) and the american heart association (aha). *Heart Rhythm* 2009;6:886-933
6. Blomstrom-Lundqvist C, Scheinman MM, Aliot EM, et al. ACC/AHA/ESC guidelines for the management of patients with supraventricular arrhythmias - executive summary. A report of the american college of cardiology/american heart association task force on practice guidelines and the european society of cardiology committee for practice guidelines (writing committee to develop guidelines for the management of patients with supraventricular arrhythmias) developed in collaboration with naspe-heart rhythm society. *Journal of the American College of Cardiology*. 2003;42:1493-1531
7. Calkins H, Kuck KH, Cappato R, et al. 2012 HRS/EHRA/ECAS expert consensus statement on catheter and surgical ablation of atrial fibrillation: Recommendations for patient selection, procedural techniques, patient management and follow-up, definitions, endpoints, and research trial design: A report of the Heart Rhythm Society (HRS) task force on catheter and surgical ablation of atrial fibrillation. *Heart Rhythm*. 2012;9:632-696.e621
8. Department of health coronary heart disease team: National service framework for coronary heart disease - chapter eight: Arrhythmias and sudden cardiac death. 2005
9. NICE. Interventional procedure guidance 168: Percutaneous radiofrequency ablation for atrial fibrillation. 2006:9
10. NICE. Interventional procedure guidance 427: Percutaneous balloon cryoablation for pulmonary vein isolation in atrial fibrillation. 2012:7
11. Campbell RWF, Charles R, Cowan JC, et al. Clinical competence in electrophysiological techniques. *Heart*. 1997;78:403-412
12. Warnes CA, Williams RG, Bashore TM, et al. ACC/AHA 2008 guidelines for the management of adults with congenital heart disease: A report of the american college of cardiology/American Heart Association task force on practice guidelines (writing committee to develop guidelines on the management of adults with congenital heart disease): Developed in collaboration with the American Society of Echocardiography, Heart Rhythm Society, International

Society for Adult Congenital Heart Disease, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *Circulation*. 2008;118:e714-e833

13. Camm AJ, Kirchhof P, Lip GYH, et al. Guidelines for the management of atrial fibrillation. *European Heart Journal*. 2010;31:2369-2429
14. Deshmukh A, Patel NJ, Pant S, et al. In-hospital complications associated with catheter ablation of atrial fibrillation in the united states between 2000 and 2010: Analysis of 93,801 procedures. *Circulation*. 2013;128:2104-2112
15. Tracy CM, Akhtar M, DiMarco JP, et al. American College of Cardiology/American Heart Association 2006 update of the clinical competence statement on invasive electrophysiology studies, catheter ablation, and cardioversion: A report of the American College of Cardiology/American Heart Association/American College of Physicians task force on clinical competence and training developed in collaboration with the Heart Rhythm Society. *Journal of the American College of Cardiology*. 2006;48:1503-1517
16. Cappato R, Calkins H, Chen SA, et al. Updated worldwide survey on the methods, efficacy, and safety of catheter ablation for human atrial fibrillation. *Circ Arrhythm Electrophysiol*. 2010;3:32-38